

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

INDEX TO VOLUME IV

New names, and the final members of new combinations, are in bold face type

```
Abies, 91, 94, 145, 146, 184, 185, 190,
                                          radiatus, 213; semiorbicularis, 76;
                                           silvaticus, 295; silvicola, 295; soli-
  261; balsamea, 145, 149, 176, 177,
  185, 187, 189, 192, 193; canadensis,
                                          tarius, 240; subnitens, 297; sub-
  184; concolor, 145; Fraseri, 177;
                                           rufescentoides, 299
  grandis, 94, 97, 145, 146; lasiocarpa,
                                        Agrimonia, 192; gryposepala, 191
  58, 65, 145, 146; magnifica, 145;
                                        Agropyron repens, 39; tenerum, 18,
  nobilis, 145, 146; pectinata, 177
                                           31, 39
                                        Agrostis alba, 18, 31; hyemalis, 9
Acer Negundo, 266; rubrum, 328
                                        Alectoria, 152; loxensis, 153; loxensis
Achlya, 320-322, 326; americana, 319;
  320; DeBaryana, 319, 320, 323, 324;
                                           atroalbicans, 155
  decorata, 325; glomerata, 325, 326;
                                        Aleuria umbrina, 221
                                        Aleuria, Artificial cultures of Asco-
  polyandra, 319, 320, 323, 324; pro-
  lifera, 319, 320, 323, 324; racemosa,
                                           bolus and, 218
  325, 326; racemosa spinosa, 325;
                                        Alopecurus, 180; pratensis, 179, 192
  racemosa stelligera, 325; spinosa,
                                        Alsine, 145
  326
                                        Althaea rosea, 18
Achyla DeBaryana Humphrey and the
                                         Amanita, 3, 231, 239; bivolvata, 241;
                                           calyptrata, 241; calyptratoides, 241;
  prolifera group, 319
Achlya glomerata sp. nov., 325
                                           calyptroderma, 241; mexicana,
                                           332; muscaria, 239; ocreata, 240;
Acolium, 152
                                           pantherinoides, 262; phalloides,
Actaea alba, 9; perennans, 9
Adoxa, 20; Moschatellina, 20, 32
                                           225; praegemmata, 262; rubes-
Aecidium, 13, 41, 195; abundans, 28;
                                           cens, 243; umbrinidisca, 262;
  auriellum, 59; elatinum, 58; monoi-
                                           virosa, 242
  cum, 59-61; Polemonii, 29; sclero-
                                         Amanitopsis, 3, 231, 239; adnata, 243;
  thecioides, 20, 59
                                           velosa, 239; volvata, 239
Aesculus glabra, 11, 12, 52, 53
                                         Ambrosia psilostachya, 23; trifida, 11,
Agaricaceae of the Pacific Coast, The
  --I, 205; II, 231; III, 294
                                         Amelanchier, 25, 26, 56, 57, 62; cana-
Agaricaceae of tropical North Amer-
                                           densis, 63; erecta, 24-26, 32, 53,
  ica, The—V, 72
                                           56, 57, 62-64; vulgaris, 25, 32, 57,
Agarics, New combinations for trop-
                                           61 - 65
                                         American mycological literature, In-
  ical, 331
                                           dex to, 160, 227, 285, 333
Agaricum, 273
Agaricus, 239, 294; Abramsii, 298;
                                         Amoeba, 322
  alneus, 213; arenicola, 81; atro-
                                         Amorpha fruticosa, 10, 12; nana, 10,
  coeruleus, 214; bivelatoides, 297;
                                           11, 13
  bivelatus, 296, 297; bulbosus, 295;
                                         Anaptychia, 126
                                         Andropogon, 17, 199, 200; furcatus,
  californicus, 294; campestris, 239,
                                           17, 31, 200, 204; glomeratus, 200;
  294, 296, 299; campestris silvicola,
  295; cerodes, 81; comptuloides,
                                           scoparius, 9, 17, 31, 200; virginicus,
  297; comptulus, 298; connatus, 216;
                                           17, 31, 200, 204
  coprinoceps, 81; crocodilinus, 300;
                                         Anemonella thalictroides, 9
  diminutivus, 298; euthugrammus,
                                         Anthoxanthum odoratum, 11
  76; flammans, 261; flavitingens,
                                         Anthurus borealis, 167
                                         Apios Apios, 10; tuberosa, 12
  298; haemorrhoidarius, 295; Her-
  barum, 244; Hillii, 298; honden-
                                         Aplanes, 322
                                         Apocynum cannabinum, 9-13, 28
  sis, 296; limonellus, 261; liratus,
  248; McMurphyi, 299; multifidus,
                                         Aquilegia canadensis, 9
  213; oinodes, 76; papularis, 80;
                                         Arabis, 59-61, 65; Drummondii, 60;
  Pattersonae, 296; pectinatus, 76;
                                           Holboellii, 10, 11, 20; retrofracta, 61
  pediades, 76; phalloides,
                                         Aragallus Lamberti, 23, 32
                                 240;
  placomyces, 295; porrigens, 215;
                                         Arenaria lateriflora, 186, 187, 193
```

```
Armillaria, 207, 211, 238; albolan-
  aripes, 211; arenicola, 212; mag-
  nivelaris, 212; mellea, 211, 266;
  nardosmia, 238; subannulata, 211
Arnica, 10, 11, 13, 20, 22, 23, 27
Aronia, 25; arbutifolia, 25, 32, 63, 65;
  nigra, 25, 32
Artemisia, 21, 32; dracunculoides, 16,
  21, 31, 32; ludoviciana, 21
Arthur, J. C., Cultures of Uredineae
in 1910, 7; in 1911, 49
Artificial cultures of Ascobolus and
  Aleuria, 218
Asclepias syriaca, 19, 31
Ascobolus, 218, 220, 221; applanatus,
  219; carbonarius, 221; laevisporus,
  219; latus, 219; magnificus, 218,
  219, 221, 222; major, 219; miniatus,
  45, 47; sarawacensis, 219
Ascobolus and Aleuria, Artificial cul-
  tures of, 218
Aspergillus, 279; flavescens, 279, 280;
  flavus, 279; fumigatus, 279; glaucus,
  279; niger, 279
Aspergillus infecting Malacosoma at
  high temperatures, 279
Aster, 7, 16, 21, 22, 144, 181, 199;
  acuminatus, 180, 181, 193; adscen-
  dens, 16, 30; Drummondii, 9, 21;
  ericoides, 19, 21, 31, 32; multiflorus,
  19, 31; Novae-Angliae, 19, 31;
  paniculatus, 10, 13, 15, 21, 28, 32,
  33; puniceus, 181, 192; Tweedyi, 16
Astragalus, 24; canadensis, 13; caro-
  linianus, 23, 32; sulphurescens, 23, 32
Atestia loxensis, 153
Atheropogon curtipendulus, 19, 31
Atriplex, 54, 55, 198; confertifolia,
  52; hastata, 54, 55, 64; patula, 179,
  198
Aurantiporellus alboluteus, 91
Auricularia auricula, 40
Auriscalpium Auriscalpium, 330
Bacidia, 126; endoleuca, 131; rubella,
  131; subgranulosa, 131
Bacterium Savastanoi, 104
Baeomyces absolutus, 132; erythrel-
  lus, 132
Bagniesiella, 70; australis, 70; Di-
antherae, 70
Banker, H. J., Type studies in the
  Hydnaceae-I. The genus Manina,
  271; II. The genus Steccherinum,
  309
Barlaea, 45, 47
Barlaeina, 45, 47
```

Berberis vulgaris, 11, 18, 31

Biatora, 126, 130-132; amorpho-

127; atrogrisea,

131;

Beta, 198

carpa,

aurigera, 127; coarctata, endocaerulea, 128; lanuginosa, 128; oncodes, 128; subgranulosa, 131; triseptata artytoides, 130 Bilimbia, 126; artytoides, 130, 131; hypnophila, 130; molybditis, 130; pallidissima, 130; radicicola, 130; sphaeroides, 130; terrestris, 130; thysanota, 130 Bierkandera adusta, 91 Black knot disease of Dianthera americana L., A, 66 Boehmeria cylindrica, 10, 17, 28 Bolbitius, 73; fragilis, 73; jalapensis, 332; mexicanus, 332; radians, 246; villipes, 73 Boletaceae of the Pacific Coast, Polyporaceae and, 91 Boletellus Ananas, 99 Boletus Lakei, 97; luteus, 98; mirabilis, 217; oregonensis, Zelleri, 217 Bolls, Botryosphaeria on cotton, 34 Boltonia asteroides, 10, 12, 20 Bombyliospora tuberculosa, 131 Botryosphaeria, 34, 36; Berengeriana, 34; fuliginosa, 34, 36 Botryosphaeria on cotton bolls, 34 Bryophagus Gloeocapsa, 127 Buellia stipitata, 139; subdisciformis, 139 Bursa Bursa-pastoris, 55 Caeoma, 30, 146, 187-189, 195; Abietiscanadensis, 188, 189, 193; conigeneum, 146; occidentale, 30 Caeomurus, 195 Calamagrostis canadensis, 18, 31 Calamovilfa longifolia, 18, 31 Calcium, 152 Calliospora, 282 Callirrhoe involucrata, 18, 31, 53 Calocera cornea, 40 Calvatia gigantea, 330 Calyptospora, 146; columnaris, 146, 177, 192 Capsella, 198 Carduus, 23; Flodmanii, 11, 22, 23, 32 Carex, 14-16, 21, 28, 54, 180, 199, 201; arctata, 9, 180, 193; aristata, 17, 31; brunnescens, 28; crinita, 180, 193; debilis Rudgei, 180, 193; deflexa, 21, 22, 32, 181, 192; festiva, 16, 30; Goodenovii, 28, 33; gracillima, 14, 201, 204; intumescens, 21, 22, 32, 180, 181, 192, 193, 199, 204; lanuginosa, 15, 30, 54, 64; pallescens, 13, 14, 30; Pennsylvanica, 7, 9; pubescens, 14, 200, 201, 204; scoparia, 15, 22, 30, 181, 182, 192, 193, 199; siccata, 16, 31; stellulata, 9; steno-

phylla, 16, 31; stipata, 181, 193; Coleosporium, 144; Vernoniae, 29, 33, stricta, 17, 31; tenella, 9; tenuis, 13, 57, 65 14, 30, 180, 201; tribuloides, 199, Collybia, 4; stipitaria, 5 Collybidium dryophilum, 164; zona-204; trichocarpa, 15, 30; trisperma, 9, 181, 193; varia, 22 Cassia chamaecrista, 11, 12 Coltricia perennis, 92 Castanea dentata, 170, 171 Comandra umbellata, 17, 31 Castilleja, 142; integra, 142 Combinations for tropical agarics, Catalpa, Polystictus versicolor as a New, 331 wound parasite of, 263 Coniophora cerebella, 223 Catillaria leptocheila, 128; rosea, Conocybe, 72, 74, 244, 247; angusticeps, 248; Bryorum, 247, 248; echinospora, 75, 332; Hypnorum, Caulophyllum thalictroides, 9 75, 247; lirata, 248; semilance-Cephalanthus occidentalis, 19, 31 Cerastium oreophilum, 58, 65 ata, 248; Sphagnorum, Cerastrum, 145 tener, 74, 247 Ceratitium, 195 Convallaria majalis, 11 Cerioniyces communis, 98, 100; mira-Coprinus, 223; Brassicae, 2; Spraguei, 5 bilis, 98, 217; oregonensis, 99, Cora pavonia, 140 217; viscidus, 99; **Zelleri,** 99, 217 Coriolellus Sepium, 92, 268, 270 Certain species of Puccinia and Uro-Coriolus abietinus, 92; nigromargimyces, Correlation between, 194 natus, 92; prolificans, 92; versicolor, Chamaedaphne calyculata, 178, 192 92; washingtonensis, 92, 217 Chanterel, 206; alectorolophoides, 206; Cornicularia loxensis, 153 behringensis, 206; bryophilus, 206; Correlation between certain species of Chantarellus, 207; floccosus, 206; Puccinia and Uromyces, 194 infundibuliformis, 206 Cortinarius, 72, 81, 243; cinnamomeus, 81; mexicanus, 81; Sintenisii, 81 Cheiranthus Menziesii, 59 Chelone glabra, 9 Corydalis aurea, 18; montanum, 55; Chenopodium, 198; album, 18, 19, 31, sempervirens, 19 Cotton bolls, Botryosphaeria on, 34 54, 55, 64, 179, 192, 198 Chestnut and oak, The large leaf spot Coursetia glandulosa, 282, 283 Crataegus, 25, 261; cerronis, 26, 32, of, 170 57, 62, 65; punctata, 25, 32; tomen-Chrysopogon avenaceus, 10 Cicuta maculata, 178, 179, 192 tosa, 24, 25, 32, 57, 63, 65 Cladonia, 132; aggregata, 132; alpes-Craterellus cornucopioides, 5 tris, 132; angustata, 132; cerato-Crepidopus, 207, 216; connatus, 216; ostreatus, 216; serotinus, 216; phylla, 132; dactylota, 132; degenerans, 132; didyma muscigena, 132; subsapidus, 216, 217 fimbriata simplex, 132; fimbriata Crepidotus, 244; calolepis, 245; fulvotomentosus, 245; Herbarum, 244; subulata, 132; Floerkeana intermedia, 132; pityrea, 132; rangiformalachias, 245; mollis, 245; puberumis, 132; sylvatica, 132. lus, 245; **submollis,** 245 Cronartium, 144; coleopsoroides, 142; Clavaria, 271, 273, 275 Quercum, 144; Quercus, 26, 33; Clematis virginiana, 11 Cleome, 54, 198; spinosa, 54, 55, 64 ribicola, 42, 104, 143 Clitocybe, 4, 330 Crouania, 45, 47; miniata, 45 Clitopilus abortivus, 6; Earlei, 332 Cryptoporus volvatus, 96 Coast, Polyporaceae and Boletaceae Cultures of Ascobolus and Aleuria, of the Pacific, 91 Artificial, 218 Coast, The Agaricaceae of the Pacific Cultures of heteroecious rusts, 175 —I, 205 ; II, 231 ; III, 294 Cultures of Uredineae in 1910, 7; in Coccocarpia pellita, 135; pellita cro-1911, 49 nia, 135 Cummings, Clara Eaton, in Jamaica, Coelocaulon divergens, 153 An enumeration of lichens collected Coenogonium Leprieurii, 127 Coker, W. C., Achlya DeBaryana by-I, 125 Cydonia vulgaris, 25, 26, 32, 56 Humphrey and the prolifera group, 319; Achlya glomerata sp. nov., 325; Dacryomyces, 40; deliquescens, 40 & Hyman, O. W., Thraustotheca Daedalea unicolor, 328 clavata, 87

Delphinium formosum, 10; scaposum, 11; tricorne, 11 Descriptions of two new species, The genus Lamprospora, with, 45 Dianthera, 66; americana, 66, 70 Dianthera americana L., A black knot disease of, 66 Diaporthe parasitica, 157 Dichodium byrsinum, 133 Dichroum, 311 Dictyolus, 206; retirugus, 206 Dictyophora duplicata, 167 Dictyuchus, 87, 322; clavatus, 87 Diplodia, 34, 36; gossypina, 34, 35 Diploschistes scruposus, 127 Dirca palustris, 9-12 Disease of Dianthera americana L., A black knot, 66 Diseases in the eastern states, Notes upon tree, 148 Distichlis, 179; maritima, 202; spicata, 12, 18, 31, 52, 54, 55, 64, 179, 192, 198, 204 Distribution, Oropogon loxensis and its North American, 152 Dodge, B. O., Artificial cultures of Ascobolus and Aleuria, 218 Doellingeria, 16; umbellata, 15 Drosophila, 294, 302; ambigua, 304; appendiculata, 302; atrofolia, 303; californica, 304; campanulata, 304; **longipes**, 303 Dryodon, 272, 276

Eastern states, Notes upon tree diseases in the, 148 Edgerton, C. W., Botryosphaeria on cotton bolls, 34 Elaeagnus angustifolia, 11, 13 Eleocharis, 203; palustris, 12 Elephantopus carolinianus, 29 Elfvingia fasciata, 105; megaloma, 96 Elymus canadensis, 11; triticoides, 39 Enumeration of lichens collected by Clara Eaton Cummings in Jamaica, An—I, 125 Epilobium, 177; angustifolium, 176, Erigeron, 7, 13, 16, 22, 199; annuus, 13 Erioderma, 135; microcarpa, 134;

Eupatorium perfoliatum, 203; serotinum, 12 Euphorbia, 202, 203; commutata, 13; corollata, 202; Cyparissias, 23, 24 Eurodium repens, 279

Euthamia, 16, 22; graminifolia, 15, 30 Evolvulus pilosus, 28, 33

Favolus, 317 Festuca confinis, 27. 33

Wrightii, 134

Flammula, 250; astragalina, 251; californica, 253; carbonaria, 165, 256; decorata, 262; echinulispora, 262; fulvella, 253; Hillii, 262; laeticolor, 262; lata, 262; ornatula, 262; pallida, 262; penetrans, 254; permollis, 262; sapinea, 254; spinulifer, 262; spumosa, 254; subcarbonaria, 262; subflavida, 262; vialis, 262; viridans, 262; viscidissima, 262 Fomes, 109; annosus, 96, 105; Earlei, 109-111, 114; fraxinophilus, 105;

juniperinus, 109, 110, 114; Laricis, 95, 97, 105; roseus, 97; texanus, 109, 110, 112, 114; ungulatus, 97

Forest trees, Notes on some western Uredineae which attack, 141

Fraser, W. P., Cultures of heteroecious rusts, 175

Fraxinus lanceolatus, 11

Friesites, 272, 276

Fungi, Illustrations of -X, 1; XI, 163; XII, 289

Fusarium, 104, 106; oxysporium, 104

Galera, 74, 247; angusticeps, 248; Besseyi, 248; Bryorum, 247; cubensis, 75; echinospora, 332; grisea, 75; Hypnorum, 75; lirata, 249; semilanceata, 248; simulans, 74; sphaerobasis, 247; Sphagnorum, 248; tener, 74; versicolor, 247 Gaura biennis, 15

Geaster juniperinus, 85, 86; marginatus, 84-86; minimus, 84-86

Geaster minimus Schw. and its relatives. Notes on Iowa saprophytes —I, 84

Gee, W. P., & Massey, A. B., Aspergillus infecting Malacosoma at high temperatures, 279

Gentiana, 203

Genus Lamprospora, with descriptions of two new species, The, 45 Genus Lasiosphaeria, The, 115 Genus Manina, The. Type studies in

the Hydnaceae-I, 271

Genus Steccherinum, The. Type studies in the Hydnaceae-II, 309 Geopetalum, 207, 214; densifolium, 215, 217; geogenium, 214; oregonense, 215, 217; porrigens, 215;

subsepticum, 215, 217 Glaux maritima, 187

Gloeophyllum hirsutum, 97

Godronia, 123; Betheli, 123; striata,

Gomphidius, 260, 294, 306, 308; nigricans, 306; oregonensis, 306; tomentosus, 307; vinicolor, 307, 308

342 Mycologia

Graves, A. H., The large leaf spot of and its North American distribuchestnut and oak, 170 tion, 152 Humaria calospora, 48 Grindelia squarrosa, 10, 19, 20, 22, 27, Hydnaceae, Type studies in the-I. Group, Achlya DeBaryana Humphrey The genus Manina, 271; II. The and the prolifera, 319 genus Steccherinum, 309 Gyalecta, 127; Gloeocapsa, 127 Hydnum, 274, 276, 317; abietinum, Gymnoconia interstitialis, 53 276; alpestre, 275; basi-asperatum, Gymnopilus, 244, 250, 261; califor-312, 313; Caput-medusae, 274, 275, nicus, 252; carbonarius, 250, 256, 277; Caput-ursi, 272, 277; conchi-257; decoratus, 250-252, 262; forme, 318; coralloides, 272-276; echinulisporus, 250, 255, 262; foedatus, 257; fulvellus, 253; Hillii, 253, 262; laeticolor, 250, croceum, 278; Daviesii, 309; decurrens, 318; dichroum, 310-312, 330; Echinus, 274; Erinaceus, 272, 275, 251, 262; latus, 257, 262; orna-277, 330; flabelliforme, 311, 312; tulus, 251, 262; pallidus, 252, glaberescens, 315; glabrescens, 312, 262; penetrans, 254; permollis, 315, 316; guaraniticum, 312, 313; hystricinum, 273, 274; Hystrix, 273, 252, 262; sapineus, 254; spinulifer, 254, 262; spumosus, 254, 274; laciniatum, 276; laeticolor, 316; 256; subcarbonarius, 256, 262; Notarisii, 275; ochraceum, 309, 310, subflavidus, 250, 252, 262; vialis, 312; Persoonianum, 312; pluma-255, 262; viridans, 250, 257, 262; rium, 317, 318; plumosum, 317; pudorium, 310-312; pusillum, 313; viscidissimus, 256, 262 Gymnosporangium, 24, 40, 41, 50, 62, ramarium, 274; ramosum, 276; 63, 106, 195, 197, 223; bermudianum, rawakense, 312, 313, 315; reniforme, 50; Betheli, 25, 32; clavariaeforme, 312; Rhois, 311, 312; rubiginosum, 24, 32, 53, 56, 64; clavipes, 24, 32; Necium Farlowii, 182, 193 cornutum, 25, 32; Davisii, 25, 32; Hydrocybe, 207; albo-umbonata, 332; durum, 61, 62, 65; effusum, 50, 62, arenicola, 208, 217; aurantia, 332; 65; Ellisii, 50; fraternum, 50; gracoccinea, 208; conica, 207; constans, 208, 217; cremicolor, 209, cilens, 63, 65; inconspicuum, 57, 64; juvenescens, 26, 62; juniperinum, 217; Earlei, 332; flavolutea, 332; 57, 65; Juniperi-virginianae, 24, hondurensis, 332; rosea, 332; sub-32; Kernianum, 62, 65; Libocedri, caespitosa, 332; subflavida, 332; 57, 65; Nelsoni, 26, 32, 61, 62, 65; subminiata, 332; troyana, 332 nidus-avis, 25, 26, 32, 56, 64; specio-Hydrophyllum capitatum, 10, 11, 27, sum, 63; tremelloides, 57, 65 29, 53; virginicum, 10, 12, 20 Hygrophorus, 207, 209; albo-umbo-Haematomma puniceum, 138 natus, 332; arenicola, aurantius, 332; coccineus, 208; Halameda, 320 Hebeloma, 72, 82, 243; Broadwayi, constans, 217; cremicolor, 217; Earlei, 332; eburneus, 209; flavo-82; cinchonense, 82; foedatum, discus, 6; flavoluteus, 332; fra-257; longicaudum, 83; subincarnatum, 83 grans, 210; hondurensis, 332; Laurae, 210; limacinus, 210; mini-Hedgcock, G. G., Notes on some western Uredineae which attack forest atus, 208; roseus, 332; rubropunctrees, 141; & Long, W. H., Pretatus, 210; subcaespitosus, 332; subflavidus, 332; subminiatus, liminary notes on three rots of juni-332; subpustulatus, 210; troyper, 109 anus, 332; variicolor, 209 Hyman, O. W., Coker, W. C. &, Hericium, 272-275; alpestre, 274, 275; Echinus, 274; grandis, 277; Hystrix, Thraustotheca clavata, 87 274; Notarisii, 274 Heteroecious rusts, Cultures of, 175 Hypholoma, 294, 302, 303, 305; am-Heterothecium, 128, 129, 131 biguum, 304; appendiculatum, 1, Hevea, 102 302; atrofolium, 303; californicum, Hibiscus militaris, 18 304; campanulatum, 304; capnoides, 305; cutifractum, 302; fasciculare, Hicoria, 121 305, 306; flocculentum, 303; lacry-High temperatures, Aspergillus infecting Malacosoma at, 279 mabundum, 302; longipes, 303, 304; sublateritium, 302 Hordeum jubatum, 39

Hypodendrum, 72, 244, 261; flam-

Howe, R. H., Jr., Oropogon loxensis

```
mans, 261; limonellum, 261;
                                           121, 123; Hystrix, 118; jamaicen-
  oregonense, 261, 262
Hypodermium, 101, 195
Hyponevris, 207, 213; alneus, 213, 214
Hypoxylon, 121; miliaceum, 121
Illustrations of fungi-X, 1; XI, 163;
  XII, 289
Impatiens, 20; aurea, 11, 20, 32
Index to American mycological litera-
  ture, 160, 227, 285, 333
Inocybe, 72, 82, 243; jamaicensis,
  82; rimosa, 4
Inonotus dryophilus, 104; texanus, 104
Iowa saprophytes, Notes on—I. Geaster
  minimus Schw. and its relatives, 84
Iris, 28; versicolor, 28
Irpiciporus, 94
Ischnoderma fuliginosum, 92
Isopyrum biternatum, 9, 11
Jamaica, An enumeration of lichens
  collected by Clara Eaton Cummings
  in-I, 125
Juneus balticus, 22, 23, 32
Juniper, Preliminary notes on three
  rots of, 109
Juniperus monosperma, 63, 65, 112, 113;
  sabinoides, 112, 113; scopulorum,
  25, 32; sibirica, 24, 25, 32, 56, 57, 64,
  65; utahensis, 56, 61, 62, 64, 65, 112,
  113; virginiana, 24-26, 32, 56, 62,
  64, 65, 85, 109, 111
Kalmia, 150; angustifolia, 150; lati-
  folia, 150
Knot disease of Dianthera americana
  L., A black, 66
Koeleria, 59, 60; cristata, 20, 31, 59
Laccaria, 207, 211; laccata, 211;
  ochropurpurea, 211
Lachnea scutellata, 38
Laciniaria punctata, 9, 10, 57; scari-
  osa, 12, 20, 29, 57; spicata, 9, 10, 12
Lactaria, 207, 289
Lactuca canadensis, 16, 31; sativa, 16,
  31; scariola, 13
Laetiporus speciosus, 92
Lamprospora, 45, 47; areolata, 48;
  tuberculata, 47
Lamprospora, with descriptions of two
  new species, The genus, 45
Large leaf spot of chestnut and oak,
  The, 170
Larix, 29, 188, 189; laricina, 29, 58,
```

188, 189

Lasiosphaeria, 115-117, 123; ambigua,

122; canescens, 118; dichroöspora, 117, 123, 124; globularis, 117, 121,

123; hirsuta, 119; hispida, 117, 119-

```
sis, 117, 122, 123; mucida, 117,
  118, 121, 124; multiseptata, 117,
  120, 124; newfieldiana, 117, 122,
  123; ovina, 118; Rhacodium, 119;
  spermoides, 121; striata, 123; stri-
  gosa, 117, 118, 123; stuppea, 117,
  119, 123; terrestris, 117, 120, 124
Lasiosphaeria, The genus, 115
Lathyrus palustris, 13
Leaf spot of chestnut and oak. The
  large, 170
Lecanora, 138; pallida, 137; subfusca,
  137; varia, 138
Lecidea, 126-131; leucoblephara, 126;
  microphyllina subgranulosa, 131
Ledum, 26, 178; groenlandicum, 26,
  33, 177, 178, 192
Lentinus lepideus, 257
Lenzites betulina, 97; sepiaria, 101
Lepargyraea canadensis, 10-13
Lepidium, 198; virginicum, 55
Lepiota, 231, 238; adnatifolia, 237;
  agricola, 332; amianthina, 237,
  238; amplifolia, 233; aspera, 238;
  asperula, 238; brunnescens, 236;
  carcharias, 237; castaneidisca,
  232; castanescens, 234; cepaes-
  tipes, 232; concentrica, 235; cre-
  tacea, 232; cristata, 232, 233; crista-
  tella, 232; eriophora, 238; fuligi-
  nescens,
            236; fulvodisca, 212;
  fumosifolia, 233; fuscosquamea,
  235; granosa, 237; granulosa, 237;
  granulosa amianthina, 238; mag-
  nispora, 237; McMurphyi, 217;
  mutata, 233; nardosmioides, 238;
  naucina, 236; petasiformis, 232;
  roseicremea,
                 217;
                       roseifolia,
  235; roseilivida, 234; rubrotincta,
  237; rubrotinctoides, 236; rugoso-
  reticulata, 237; Sequoiarum, 233;
  subfelina, 234; subnivosa, 231
Leptogium bullatum, 133; chlorome-
  lum, 133; foveolatum, 134; lacerum,
  133; marginellum, 133; myochroum
  tomentosum, 134; phyllocarpum,
  133; punctulatum, 133; saturninum,
  134; tremelloides, 134; tremelloides
  impresso-punctatum, 134
Leptonia atrosquamosa, 332; cin-
  chonensis,
              332; Earlei,
  mexicana, 332
Leptoniella atrosquamosa, 332; cin-
  chonensis, 332; Earlei, 332; mexi-
Leptospora, 115, 117; ovina, 118;
  spermoides, 121; strigosa, 118
Leucomyces, 239; mexicanus, 332
Lewis, I. M., A black knot disease of
  Dianthera americana L., 66
```

Libocedrus decurrens, 57, 65

Lichens collected by Clara Eaton Cummings in Jamaica, An enumeration of-I, 125 Limacella, 207, 212; agricola, 332; fulvodisca, 212; McMurphyi, 213, 217; roseicremea, 212, 217 Literature, Index to American mycological, 160, 227, 285 333 Lithospermum canescens, 10, 11 Lobaria, 135; corrosa, 135; erosa, 136; pallida, 135; peltigera, 135; quercizans, 136 Locellina, 244, 262; californica, 262; stercoraria, 262 Long, W. H., Two new species of rusts, 282 Long, W. H., Hedgcock, G. G. &, Preliminary notes on three rots of juniper, 109 Lopadium amaurum, 131; leucoxanthum, 131 Lophodermium nervisequum, 149 Lycopus americanus, 17, 31, 54, 64 Lygodesmia juncea, 58, 65 Lysimachia quadrifolia, 9, 10; terrestris, 9, 12, 13, 22, 28 Macbride, T. H., Notes on Iowa saprophytes-I. Geaster minimus Schw. and its relatives, 84 Macrocalyx Nyctelea, 11, 28 Macrophoma, 34, 103 Mahonia Aquifolium, 11 Malacosoma americana, 279, 280 Malacosoma at high temperatures, Aspergillus infecting, 279 Malus, 56; coronaria, 26, 56, 63, 64; Malus, 24, 32, 63 Malvastrum coccineum, 52 Manina, 271-276, 278; Caput-ursi, 277; coralloides, 276, 330; cordiformis, 272, 275, 277, 330; flagellum, 276; Schiedermayeri, 277 Manina, The genus. Type studies in the Hydnaceae—I, 271 Marasmius, 4, 5, 157; magnisporus, 166; oreades, 224; Vaillantii, 167; viticola, 167 Martela, 273 Martella, 273-275; Echinus, 273, 274 Massey, A. B., Gee, W. P., &, Aspergillus infecting Malacosoma at high temperatures, 279 Medicago, 24, 56; sativa, 13, 24, 32, 56, 64 Medusina, 272, 275; coralloides, 276; patula, 272, 275, 277 Megalospora Cummingsiae, 129; jamaicensis, 129; sulphurata, 129; versicolor, 129

Melampsora, 30, 146, 188, 189; albertensis, 29, 33, 58, 65; arctica, 187, 193; Bigelowii, 189; Medusae, 188, 189, 193 Melampsorella elatina, 58, 65, 145 Melampsoropsis, 175; abietina, 26, 33, 178, 192; Cassandrae, 178, 192; ledicola, 177, 192; Pyrolae, 183, 184, Melanoleuca jalapensis, 332; jamaicensis, 332; subisabellina, 332 Melanops Quercuum, 34 Meriolix, 15; serrulata, 9, 15, 30, 54 Merisma, 274 Merulius lacrymans, 223, 327 Metasphaeria sublanosa, 118 Microsphaera quercina, 223 Microphiale lutea, 127 Mimulus ringens, 10, 12 Monochaetia, 172; Desmazierii, 104, 170-174, 330, 331; monochaeta, 172; pachyspora, 170, 331 Monolepis Nuttalliana, 18, 19, 54 Mucor, 88 Muhlenbergia gracilis, 18; racemosa, Murrill, W. A., Illustrations of fungi -X, 1; XI, 163; XII, 289; New combinations for tropical agaries, 331; Polyporaceae and Boletaceae of the Pacific Coast, 91; The Agaricaceae of the Pacific Coast-I, 205; II, 231; III, 294; The Agaricaceae of tropical North America-V, 72 Mycelophagus Castaneae, 223 Mycena, 72, 73; fragilis, 73; jalapensis, 73, 332; mexicana, 73, 332; praedecurrens, 165; vexans, 168; villipes, 73; vulgaris, 165 Mycological literature, Index to American, 160, 227, 285, 333 Myrica cerifera, 9, 10, 11, 28 Myxosporium acerinum, 151

Naematelia encephala, 40; nucleata, 40
Napaea dioica, 18
Naucoria, 72, 74, 75, 81, 243; corticola, 77; cyathicola, 77; Earlei, 77; euthugrammus, 76; hepaticicola, 78; jalapensis, 77; montana, 78; oinodes, 76; paludosa, 249; pectinata, 76; pediades, 76; pellucida, 78; Sacchari, 79; semiorbicularis, 76; sideroides, 80; spinulifer, 79; subvelosa, 164; tepeitensis, 79; Underwoodii, 80; xuchilensis, 80
Necium Farlowii, 182, 193
Nectria cinnabarina, 266
New combinations for tropical agarics, 331

```
244; involutus, 244; ligneus, 73;
New species of rusts, Two, 282
                                           panuoides, 73
New species, The genus Lampro-
  spora, with descriptions of two, 45
                                         Peltigera canina, 137; polydactyla,
News and notes, 37, 101, 157, 223, 327
                                           137
                                         Pentstemon alpinus, 17, 31; barbatus,
Nigredo, 195, 223
North America, The Agaricaceae of
                                           17; hirsutus, 17, 31
  tropical-V, 72
                                         Peridermium, 50, 141-146, 184, 189,
North American distribution, Oro-
                                           190, 195, 282, 284; abietinum, 178;
  pogon loxensis and its, 152
                                           balsameum, 189-191; carneum, 29,
Notes, News and, 37, 101, 157, 223,
                                           57; Cerebrum, 26, 143, 144; colo-
                                           radense, 145; Conorum-Piceae, 146,
  327
Notes on Iowa saprophytes—I. Geaster
                                           183, 184, 193; consimile, 178; de-
  minimus Schw. and its relatives, 84
                                           colorans, 178; delicatulum, 282, 284; elatinum, 145; filamentosum,
Notes on some western Uredineae
  which attack forest trees, 141
                                           141-144; fructigenum, 150, 185,
                                           186; Harknesii, 143, 144; incon-
Notes on three rots of juniper, Pre-
                                           spicuum, 283; montanum, 144,
 liminary, 109
Notes upon tree diseases in the east-
                                           284; Peckii, 186, 193; pseudo-
  ern states, 148
                                           balsameum, 146; stalactiforme, 142;
                                           Strobi, 42, 143
Oak, The large leaf spot of chestnut
                                         Pertusaria cryptocarpa, 137; leiopla-
                                           cella, 137; tuberculifera, 137; velata,
  and, 170
Ocellularia actinotum, 126; anamor-
                                           137
  phum, 126; auberianoides, 126;
                                         Pestalozzia, 172, 331; monochaeta, 172
  clandestina, 127; terebrata, 127
                                         Petalostemon purpureus, 10-12
Odontia fimbriata, 330
                                         Phacelia heterophylla, 10
                                         Phaeolus sistotremoides, 93
Oidium, 223
Omphalia, 165; Austinii, 165; Cam-
                                         Phalaris arundinacea, 11
  panella, 168, 250
                                         Phialocybe, 72
Omphalopsis Campanella, 168
                                         Philadelphus, 63, 64; coronarius, 62,
Onagra, 15; biennis, 15, 16, 30, 54,
                                           63, 65
  64; pallida, 53
                                         Phlebia, 317; hydnoides, 278
Onoclea, 190; sensibilis, 190, 191
                                         Phlox, 29; divaricata, 12, 29
Onosmodium hispidissimum, 11; occi-
                                         Pholiota, 72, 244, 258; aeruginosa,
                                           257; albivelata, 260; anomala,
  dentale, 10
Orchis spectabilis, 9
                                           259; autumnalis, 330; candicans,
Oropogon, 152, 153; loxensis, 152-154
                                           259; marginata, 258; McMurphyi,
Oropogon loxensis and its North
                                           260; oregonensis, 262; praecox,
  American distribution, 152
                                           259; subnigra, 258; unicolor, 258;
Orton, C. R., Correlation between cer-
                                           ventricosa, 260; washingtonen-
  tain species of Puccinia and Uro-
                                           sis, 259
  myces, 194
                                         Pholiotina, 72
Otomycosis aspergillina, 279
                                         Phoma, 106; piciena, 151; pigmen-
Oxygraphis Cymbalaria, 54, 64
                                           tivora, 38
Oxytropis, 23; Lamberti, 23
                                         Phragmidium, 194, 195, 197
                                         Phragmites communis, 54, 64
Pacific Coast, Polyporaceae and Bole-
                                         Phragmopyxis, 282
taceae of the, 91
Pacific Coast, The Agaricaceae of the—I, 205; II, 231; III, 294
                                         Phryma leptostachya, 9
                                         Phyllopsora furfuracea, 131; parvi-
                                           folia, 132
Panicum virgatum, 12, 202, 204
                                         Physalis, 11
Pannaria, 134; leucosticta, 134; molybdea, 135; rubiginosa, 134
                                         Physcia, 126; comosa, 140; hypo-
                                           leuca, 140; leucomela, 140
Panus stypticus, 77
                                         Physcidia Wrightii, 138
Parasite of catalpa, Polystictus versi-
                                         Physma byrsaeum, 133
  color as a wound, 263
                                         Picea, 26, 178, 183, 184; canadensis,
Parmelia cetrata ciliosa, 138; cetrata
                                           177, 178, 183, 192, 193; Engelmanni,
                                           145, 146; excelsa, 151; Mariana, 26,
  subsidiosa, 138; laevigata, 138;
  perforata, 138
                                           33, 178, 183, 184, 192, 193; Parry-
Parmeliella tryptophylla, 134
                                           ana, 145; rubra, 151, 178, 192;
```

sitchensis, 145

Paxillus, 72, 243, 244; atrotomentosus,

96

Pilocarpon tricholoma, 126 Psilocybe, 330 Puccinellia airoides, 54, 64 Pinus, 42; chihuahuana, 146; contorta, 142-144; Jeffreyi, 143; pon-Puccinia, 9, 11, 14, 21, 22, 55, 56, 180, derosa, 141, 143; radiata, 143; 181, 194-197, 199, 200, 203, 204; rigida, 282; sabiniana, 143; taeda, Absinthii, 21, 32; albiperidia, 13 29, 33, 57, 65; virginiana, 26, 33, (see page 49), 14, 15, 30 (see page 282, 284 49), 49, 180, 193, 200, 201; amphi-Pisum sativum, 13 gena, 18, 31; Andropogonis, 7, 17, Pleurage, 116; albicans, 116; lutea, 31; angustata, 7, 17, 31, 54, 64; angustatoides, 203; Anthoxanthi, Pleuropus abortivus, 6; Earlei, 332 11; argentata, 20, 32; Caricis, 7, 17, 31; Caricis-Asteris, 16, 28, 30, 181, Pleurotus, 4, 207, 217; candidissimus, 193, 199, 204; Caricis-erigerontis, 215; corticatus, 217; densifolius, 217; dryinus, 217; geogenius, 3; 199; Caricis-Solidaginis, 15, 30, 181, oregonensis, 217; ostreatus, 216; 193, 199; cinerea, 54, 64; Crandallii, sapidus, 159, 216; septicus, 215; serotinoides, 216; serotinus, 216; 27, 33; Distichlidis, 11, 202, 204; Eatoniae, 61; Eleocharidis, 203; Ellisiana, 7, 9, 199, 200, 204; emacusubareolatus, 217; subsapidus, 217; subsepticus, 217 lata, 7; fraxinata, 7; Gentianae, Plicaria echinospora, 221 203; graminis, 39, 107; Grossu-Plowrightia, 70 lariae, 14; see page 49 for error; heterantha, 203; Impatientis, 7; Jamesiana, 19, 31; Lithospermi, 28, Pluteolus, 72, 74, 244, 246, 247; californicus, 246; callistus, 247; expansus, 247; luteus, 246, 247; 33; Lygodesmiae, 58, 65; malvaparvulus, 246; stramineus, 246; cearum, 329; monoica, 61, 65; tropicalis, 74 montanensis, 11; Muhlenbergiae, Pluteus stercorarius, 262 18, 31; opaca, 203; Opizii, 16, 31; Podiosoma, 195 Panici, 202; Pammellii, 202-204; Polemonium, 29; reptans, 10, 12, 28, Peckii, 7, 15, 30, 54, 64, 181; permi-29, 33 nuta, 9; perplexans, 179, 192; Polygala Senega, 12, 13 Phragmites, 54, 64; poculiformis, 7, 18, 31; pustulata, 7, 17, 31; quadriporula, 28, 33; Rhamni, 11, Polygonatum commutatum, 11 Polyporaceae and Boletaceae of the Pacific Coast, or 18, 31; Ruelliae-Bourgaei, 203; Polyporus carbonarius, 217; Car-Seymouriana, 19, 31; Schedonneus, 109; cutifractus, 217; elenardi, 10, 53; Stipae, 19, 31, 59; gans, 93; juniperinus, 109, 110; subnitens, 18, 31, 52, 54-56, 64, 198, oregonensis, 217; perdelicatus, 204; tosta, 10, 52; Triseti, 59; uni-217; Pseudotsugae, 217; sensiporula, 201, 204; universalis, 16, bilis, 217; substipitatus, 217 31; vexans, 7; virgata, 10 Polystictus abietinus, 113; versicolor, Puccinia and Uromyces, Correlation 263, 264, 266-270; washingtonbetween certain species of, 194 ensis, 217 Pucciniastrum agrimoniae, 191; arc-Polystictus versicolor as a wound tium, 190, 191; minimum, 184, 186, parasite of catalpa, 263 193; Myrtilli, 186; pustulatum, 176, Populus, 30, 189; acuminata, 147; 177, 192 angustifolia, 147; balsamifera, 147; Pulicaria, 23 grandidentata, 147, 188, 193; tremu-Pyrola, 183, 184; americana, 184, 193; loides, 29, 33, 58, 65, 147; trichoelliptica, 184, 193 carpa, 147 Pyropolyporus Earlei, 111; Everhartii, Poria, 113; vaporaria, 223 104; igniarius, 97, 104; juniperinus, Porodaedalea Pini, 97, 105 110; texanus, 112 Preliminary notes on three rots of Pyrus communis, 25, 63 juniper, 109 Pyxine picta, 139 Prolifera group, Achlya DeBaryana Humphrey and the, 319 Quercus Ilex, 172; nigra, 331; Prinus, Psathyra, 81 103; rubra, 26, 33, 170, 172, 331 Psathyrella, 303 Pseudotsuga, 29, 30, 58, 261, 330; Ramalina denticulata, 138; linearis, mucronata, 29, 33, 58, 65; taxifolia,

138; usneoides, 138

Sarcobatus, 198; vermiculatus, 54 Ranunculus abortivus, 61; acris, 179, Schedonnardus paniculatus, 10, 53 Razoumofskya cryptopoda, 142 Schizophyllum, 213; commune, 213; Relatives, Geaster minimus Schw. and Egelingianum, 214; exiguum, 214; fasciatum, 213; flabellare, 214; mexicanum, 214; multifidum digiits. Notes on Iowa saprophytes-Resupinatus, 207, 214; atrocoerutatum, 214; pavonium, 214; pusillum, 214; umbrinum, 213, 214 leus, 214 Rhamnus, 11; alnifolia, 11, 18, 31; Schizophyllus, 213 Scirpus atrovirens, 17, 31; campestris cathartica, 11 Rhododendron, 150; arborescens, 150; paludosus, 178, 192; cyperinus, 54, 64; fluviatilis, 179; paludosus, 178; calendulaceum, 150; canadense, 150; validus, 179 canescens, 150; catawbiense, 150; Scutiger oregonensis, 93, 217; renudiflorum, 150; maximum, 150; viscosum, 150 tipes, 93 Rhodora, 184, 185; canadense, 184, Seaver, F. J., The genus Lamprospora, with descriptions of two new spe-193 cies, 45; The genus Lasiosphaeria, Rhyncospora, 203 Ribes, 14, 42, 180, 200, 201; Cynosbati, 9, 13, 30, 58, 180, 200; floridum, 28; Senecio Douglasii, 9, 11; lugens, 11, oxyacanthoides, 180, 193; prostra-20, 28, 31 tum, 180, 193 Septoria ochroleuca, 104, 170 Ricasolia subdissecta, 136 Sidalcea oregana, 10, 22, 53 Riddle, L. W., An enumeration of Sitanion longifolium, 18, 31 lichens collected by Clara Eaton Smilax hispida, 18, 31 Cummings in Jamaica—I, 125 Solidago, 7, 16, 22, 181, 199; bicolor. Rinodina Conradi, 139; exigua, 139; 181, 192; caesia, 9; canadensis, 9, sophodes exigua, 139 13, 15, 19, 21, 31; graminifolia, 181, Roestelia hyalina, 50; transformans, 192, 193; lanceolata, 15; nemoralis, 15, 21; rigida, 10; rugosa, 15, 21, Rostkovites granulatus, 100 32, 181, 193 Some western Uredineae which at-Rots of juniper, Preliminary notes on tack forest trees, Notes on, 141 three, 109 Sorbus, 25; americana, 25, 32, 57, 65; Rubus, 190, 191; allegheniensis, 53 Rudbeckia laciniata, 9, 12, 13, 28; triaucuparia, 57, 63 Spartina, 11, 12, 186, 201; cynosuroides, 12, 29; glabra alterniflora, loba, 9–11 Ruellia, 203 Rumex, 196; crispus, 54, 64 186, 187, 193; gracilis, 201, 202, Russula, 207, 289, 291; foetens, 292; emetica, 290; Mariae, 290, 293; 204; Michauxiana, 12, 19, 29, 31, 33, 186, 187, 193, 201; patens, 12, 186, obscura, 291; rubriochracea, 293; 187, 193 sericeonitens, 290; stricta, 166; Spaulding, Perley, Notes upon tree sulcatipes, 291; uncialis, 292 diseases in the eastern states, 148 Species of Puccinia and Uromyces. Rusts, Cultures of heteroecious, 175 Rusts, Two new species of, 282 Correlation between certain, 194 Species of rusts, Two new, 282 Ryssospora, 72 Species, The genus Lamprospora, with descriptions of two new, 45 Salicornia, 179; europaea, 179, 187, Spergularia canadensis, 179, 186, 187, Salix, 147; amygdaloides, 147; Beb-Sphaeria canescens, 118; emergens, biana, 147; cordata lutea, 147; cordata Mackenziana, 147; discolor, 119; globularis, 121; hirsuta, 119; 188, 193; fluviatilis, 147; laevigata, hispida, 119; mucida, 118; mutabilis, 118; orthotricha, 119; ovina, 147; lasiandra, 147; lasiandra cau-115, 117, 118; Rhacodium, 119, 121; data, 147; lucida, 147; nigra, 147; Nuttallii, 147; rostrata, 188; sessispermoides, 121; strigosa, 118; subfolia, 147 lanosa, 118; terrestris, 120 Sambucus racemosa, 180 Sphaerophorus compressus, 126 Saprolegnia asterophora, 326 Sphaeropsis, 34, 36; tumefaciens, 329 Spongipellis sensibilis, 93, 217 Saprophytes, Notes on Iowa — I. Geaster minimus Schw. and its rela-Sporobolus asperifolius, 10, 52; neg-

lectus, 13

tives, 84

Thelephora, 317; multifida, 317

Thraustotheca clavata, 87

notes on, 109 Thuya, 95; plicata, 92

Thelotrema, 126; concretum, 127

Thraustotheca, 87, 88; clavata, 87

Tissa canadensis, 11, 12, 19, 21, 55

Three rots of juniper, Preliminary

Steecherinum, 309, 317; adustulum, Tonina janeirense, 131 basi-badium, Trametes Pini, 145; Sepium, 268 314; chroum, 310, 311; laeticolor, Tree diseases in the eastern states, 316, 318; Morgani, 315, 316; mul-Notes upon, 148 tifidum, 317, 318; ochraceum, 310-Trees, Notes on some western Ure-312, 317, 318; Peckii, 314, 315; dineae which attack forest, 141 pusillum, 313; quercinum, 277; Tremella, 40 rawakense, 312, 313, 315; Rhois, Tricella, 282; acuminata, 282 310, 311, 314 Tricholoma jalapensis, 332; jamai-Steccherinum, Thegenus. Type studies censis, 332; subisabellina, 332 in the Hydnaceae-II, 309 Trifolium, 24; medium, 24, 56; pra-Steironema, 12, 202; ciliatum, 12, 29, tense, 13, 24, 56; repens, 24, 56 Trillium cernuum, 11 Stellaria graminea, 187; media, 187 Triphragmium, 195 Trisetum, 59-61; majus, 11, 59, 65; Stereocaulon cornutum, 133; pityrizans, 133; ramulosum, 133 subspicatum, 59, 60, 65 Tropical agaries, New combinations Stereum purpureum, 105 Stevens, N. E., Polystictus versicolor for, 331 as a wound parasite of catalpa, 263 Tropical North America, The Agarica-Sticta, 135; aurata, 136; crocata, 136; ceae of-V, 72 crocata leucosticta, 136; damae-Tsuga, 184, 185, 189, 191; canadensis, cornis, 136; damaecornis elongato-150, 176, 177, 182, 184-186, 188, laciniata, 136; damaecornis sinu-189, 191, 193; ciniastrum minimum, osa, 136; dissecta, 135; dissecta corrosa, 135; erosa, 136; Fendleri, 185; heterophylla, 95; Pattoniana, 119 135; Mougeotiana, 136; peltigera, Tubaria, 244, 249, 259; brevipes, 250; 136; quercizans, 137; tomentosa, contraria, 249; deformata, 249; 137; Weigelii, 137 Eucalypti, 249; furfuracea, 249; in-Stipa, 19, 31, 59, 60; spartea, 19, 31; quilina, 249; pallescens, 249; pelluviridula, 59 cida, 79; tenuis, 249 Strobilomyces strobilaceus, 99 Two new species of rusts, 282 Stropharia, 294, 300, 305; aeruginosa, 300; bilamellata, 302; longistri-Two new species, The genus Lamprospora, with descriptions of, 45 Type studies in the Hydnaceae-I. The ata, 301; magnivelaris, 301; semiglobata, 2, 300, 301; semigloboides, genus Manina, 271; II. The genus 301; stercoraria, 300, 301 Steccherinum, 309 Studies in the Hydnaceae, Type-I. Tyromyces caesius, 94; carbonarius, The genus Manina, 271; II. The 94, 217; chioneus, 94; cutifractus, genus Steccherinum, 309 94, 217; guttulatus, 95; perdelicatus, 95, 217; Pseudotsugae, 95, Suedia, 179; maritima, 179 Suillellus luridus, 100, 163 217; semipileatus, 96; semisupinus, Symphoricarpos oreophilus, 28; pau-96; substipitatus, 96, 217 ciflorus, 10; racemosus, 10-12, 20, Urceolaria, 127 27, 33 Uredineae in 1910, Cultures of, 7 Tachaphantium, 40 Uredineae in 1911, Cultures of, 49 Tapinia, 72, 243, 244; lignea, 73; Uredineae which attack forest trees, panuoides, 244 Notes on some western, 141 Teloschistes flavicans, 139 Uredinopsis, 190, 191; mirabilis, 190, Temperatures, Aspergillus infecting 191; Phegopteridis, 191; Osmundae, Malacosoma at high, 279 191 Thalictrum dioicum, 9; polygamum, 10 Uredo, 194, 195, 197; Bigelowii, 147; Thalloidima, 131 Fabae, 194; Medusae, 147; Oxy-

tropi, 23
Uromyces, 12, 14, 21, 22, 55, 56, 179, 186, 194–197, 199, 201, 203, 204, 223; Astragali, 23, 32; acuminatus, 29, 33, 201, 202, 204; caricina, 22, 199; cucullatus, 203; Eleocharidis, 12, 203; graminicola, 12, 202, 204; Junci, 22, 32; Medicaginis, 24, 32,

56, 64; Peckianus, 12, 55, 56, 64, 179, 192, 198, 204; pedatatus, 200, 204; perigynius, 21, 22, 32, 181, 192, 199, 204; plumbarius, 203; Rhyncosporae, 203; Ruelliae, 203; Scirpi, 178, 179, 192; Spartinae, 12, 186, 187, 193; speciosus, 203; Sporoboli, 13; Solidagini-Caricis, 22; uniporulus, 14, 201, 204 Uromyces, Correlation between certain species of Puccinia and, 194 Urtica gracilis, 17, 28, 31 Usnea angulata, 138; ceratina, 138; dasypoga, 139; florida, 139; laevis, 139; plicata, 139 Ustilago esculenta, 104 Uvularia grandiflora, 10, 11

Vaccinium, 146; canadense, 186; pennsylvanicum, 177, 186, 192
Vaginata, 231, 239; farinosa, 3; vaginata, 239; velosa, 239
Vagnera stellata, 11
Venenarius, 231, 239; bivolvatus, 241; calyptratoides, 241; calyptratus, 241, 242; mexicanus, 332; muscarius, 240, 243; ocreatus, 240;

pantherinoides, 242, 262; phalloides, 240; praegemmatus, 243, 262; solitarius, 240; umbrinidiscus, 242, 262
Vernonia, 29; crinita, 29, 33; gigantea, 57, 65
Verticillium, 104
Viburnum, 318
Viola, 199, 200; septentrionalis, 11
Viorna Douglasii, 11
Vitis, 34
Volvaria Bakeri, 332; cubensis, 332; Earlei, 332; jamaicensis, 332
Volvariopsis Bakeri, 332; cubensis,

Western Uredineae which attack forest trees, Notes on some, 141 Wound parasite of catalpa, Polystictus versicolor as a, 263

332; Earlei, 332; jamaicensis, 332

Xanthoxylum americanum, 10, 52, 53

Zexmenia, 203 Zizania latifolia, 104